Identification of an RNase that preferentially cleaves A/G nucleotides

Jumin Xie^{1, 2} Zhen Chen¹ Xueyan Zhang^{1, 2} Honghe Chen¹ Wuxiang Guan ^{1*}

¹Center for Emerging Infectious Diseases, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, Hubei, China.

² University of Chinese Academy of Sciences, Beijing, 100049, China.

^{*}Correspondence should be addressed to W.G. (guanwx2000@hotmail.com).

Supplementary table S1. 3 independent mass spectrum analysis of hTRIR.

	Accession	Description	ΣCoverage	Σ# Proteins	Σ# Unique Peptides	# AAs	MW [kDa]
	Q9BQ61	Uncharacterized protein C19orf43 OS=Homo sapiens	7.95	1	1	176	18.4
1		GN=C19orf43 PE=1 SV=1 - [CS043_HUMAN]					
		Campana	# DCM-	#	# Protein	# Missed	
		Sequence	# PSMs	Proteins	Groups	Cleavages	
		EAPGPAGGGGGSR	2	1	1	0	
	Q9BQ61	Uncharacterized protein C19orf43 OS=Homo sapiens	22.73	1	3	176	18.4
2		GN=C19orf43 PE=1 SV=1 - [CS043_HUMAN]					
		Sequence	# PSMs	#	# Protein	# Missed	
		Sequence	# F3145	Proteins	Groups	Cleavages	
		QRQEEPPPGPQRPDQSAAAAGPGDPKR	3	1	1	2	
		QRQEEPPPGPQRPDQSAAAAGPGDPKR	3	1	1	2	
		KGGPGSTLSFVGK	2	1	1	1	
		GGPGSTLSFVGK	2	1	1	0	
	Q9BQ61	Uncharacterized protein C19orf43 OS=Homo sapiens	54.55	1	9	176	18.4
3		GN=C19orf43 PE=1 SV=1 - [CS043_HUMAN]					

Supplementary figure S1.




Figure S1. hTRIR is an exoribonuclease. (A) Chemically synthesized RNA oligos with methylated group at either 5' end or 3' end were

shown. (B) Synthesized 1A and (C) A1 were digested by hTRIRD for different time as indicated. Low molecular ladder of digested 1A and A1 were detected after 5 min incubation and marked with *.